Collection no. 189

OLYMPIC

GLAMIS

Key, Balderson, + Freeman

DUPLICATE

CARBON BOOK

(Octavo Size)

Gove Peninsula, N.T.

100 Leaves in Duplicate

15 – 24.V.1975

NUMBERED

Flexiflat

THERMO–PLASTIC BOUND

Collection no. 189 — Cove Peninsula, N.T.

Key, Balderson, & Freeman

15–24.V.1975

All odometer readings

15/5 Plants det. M. Bazaroides, R. Hooglandi (Hibbertia) in Rm.

11020.5 Nhulunbuy motel. Took road to airport.

11024.8 Gentle undulations. Brown Rectitropis sp. soil with laterite gravel. Open enc. forest. Two spp. encs (spaces) to 50ft, sp. 12, Goniaceae vocans, Austrocactus bifalcis, Macrotona sp. 2, Goniaceae sp. 3, Heteroptenis obscurella, Macrazolota sp. 2, Xanterriana concolor, Genus nov. 105 sp. 1 one a woodwood, both rough to apex. (j.), Erythropomala amaena (j.), Ndecrocarpa onglobata, Acacia spp (3, phyllod.) to 5ft, Calythrix Gastrinargus muricatus, Abbeckia rupescens, Bundinja enc. to 5ft, palms to 8ft, clado-de-type sp. P70, Spectrifoma transversa, Hostella sp. P208. papillonaceous spiny shrub (as Petrocoda camp) to 4ft, Hibbertia dealbata (DC.) Benth., H. lepidota D.C. Hibbertia spp. (2) Boronia ?artemisiifolia F. Muell. mainly to 2ft, a ~~forest~~ belt to 18m (spec.) Grasses along road.

16/5

11029.1 Nhulunbuy motel.

Mt. Saunders. 70m. Flat-topped hill Goniaceae sp. 3, Stenocatantops angustifrons, with massive laterite. Rough-barked S. vitripennis, Xanterriana concolor, Caledia (2) enc, slender, to 25ft, with Acacia spp. captiva, Diolopus thalassinus. Heteroptenis (phyllodineus) to 12ft, Miscellaneous obscurella, Tolgadia inferna, Goniaca vocans, broad-leafed shrubs to 4ft & scramblers. Acrida conica, Austracris guttulosa, Rectitropis Dec. palms to 6ft, mixed grasses to sp., Bundinja sp. P70 3ft. Tract. Moraine on Acacia spp.

11036.9 Motel (miles wasted)

Erythrophonala amaena, Genus nov. 28 sp. 1, Equivalvidea sp. 3, Xanthetraria conc. sp., Austraeris basalio, Macrozelota sp. 2, Goniura vocans, Daperia accola, Adlappa erythroptera, Abbetia rufescens, Bourdiuja sp. P70, Spectrifonna Pandanus. Hailsoderma

11041.2 = 11024.8. Daperia associated with

042.0. Road off to left.

043.3. Road off to left.

047.0. Road off to left to Gun Club.

048.8. Road off to left to Ginkala + to right.

049.7. Crossed conveyor belt. Mine on left.

052.1. Airport entrance on right.

054.9. Sharp bend to R.

058.9. Flat adjacent to small creek (Latram River).

Stenocatantops ultripennis, Calidia capstiva, Brown soil. E. ? Tetrodonta, occ. E. Rectitropis sp.

Acacia to 70ft. Acacia spp. phyllod. (2) to 20ft, occ. palms, occ. pandanus occ. Hibbertia to 1ft, grasses abundant to 5ft.

061.1. Track off to L.

065.2. Spur on L.

066.0. Dhupuma College.

066.1. Turned left

067.4. Edge of laterite scarp with view over to sand dunes + sea. Flat, pale

Heteropternis obscurella, Stenocatantops
brown soil with lateritic gravel. E.
nitripennis, Acrida open conch.
? tetrodonta to 30 ft, with Acacia spp. h
to 15ft. + Erythrophleum regrowth to 5ft.
A. sp. 2, Gonaea volans, Caledia captiva,
Patches of melaleuca spp. to 15ft + regrowth.
Tall (to 4ft) + short grasses, occ.
Spectriforma brunneriana, Hastilla sp. P208
Hibbertia sp. to 1ft + occ. other shrubs
to 3ft. Return.

068.7 Back on main road. Turned L.

069.9. Fork. Kept R.

071.2. End of road. Scarp overlooking
Adappa erythropteta, Tolgadia inferna,
sand-dunes + sea. E. tetrodonta to
Spectriforma transversa
30 ft, with Acacia spp. & other small
trees to 15ft. Erythrophleum regrowth to
4ft, grass, occ. Hibbertia + forbs.
Return.

17/5.
am.
Beach dunes & adjacent swamp SE of
hotel. (1) Broad-leafed low trees + shrubs
(Celastraceae), Celtis philippensis Blanco (Ulmaceae),
on white sand at steep fall to beach.
Diospiros calycantha O. Schwartz (Ebenaceae),
Valanga (3 spp.) basking on these at
Drypetes australasica (Muell. Arg.) Pax & Hoffm.
0830 – 0900 hr, easily caught by hand or
Valanga meleager, V. sp. 1, V. irregularis
net. By 0930 more active: diving

Elaeodendron melanocarpum F. Muell.

into vegetation. (2) Nearer to sea, on Austrapicis basalis, Genus nov. 70 sp. 1, Daperia white sand flats. Casuarina sp. to accola
25ft with coarse tussock grasses, forbs, occ. Pandanus + broad-leaf shrubs, trailing Ipomoea & other trailers, Stenocatantops angustifrons S. vitripennis, Peyerostictus variatus, Aiolopus, Gonracridea sp. & Gen. nov. 70(?). Daperia associates with Pandanus.

(3) Samphire to 3ft on pale sandy mud, Arthrocnemum sp.
Genus nov. 67 sp. 1
wet, with numerous crabs, passing into mangroves. Living estuary.

(4) Swampy area with Melaleuca Gonracridea sp. 3, Canterriana concolor, Genus sp. to 20ft, Pandanus to 15ft & nov. 70 sp. 1, Tolgadia bivittata, Daperia regrowth, occ. Phyllodineous Acacia accola, Rectitropis sp., Acrida acuta, conica, sp. to 10ft, mixed grasses & sedges Caledia captiva, patch of Blundunga sp. P70, Hastella to 4ft — beach. Tolgadia sp. P208 Daperia sp., sp., Stenocatantops vitripennis. Parabiine on Acacia.

pan. (5) Billabong margin & dune complex Gonracridea sp. 3, Genus nov. 70 sp. 1, Atractomorpha lutum & N of hotel. Melaleuca to 40ft similis, Tolgadia inferna, Acrida conica, along billabong, with hippophilous Rectitropis sp., Valanga sp. 1, Xyrchtria sp. 1, grasses, sedges, & forbs & rush & Aiolopus thalassinus, Paratettix nigrocens

mud along water's edge. Patches of dense dune trees + shrubs, including Ficus spp., becoming larger + more frequent northwards. Seaward dunes with Casuarina to 30ft. including Spinifex sp., Grasses + forbs, much bare sand. Tetragidia + Abrotanella fringing water.

18/5.

11106.4 Hotel.

109.7 Sewage treatment works. Return.

110.3 Margin of sandy swamp. A bloodwood Osaga japonica, Heteropteris obscurella, + very broad oblong-leafed one, with Rhetinopis sp., Macrozamia Conglobata, Zamiaceae Melaleuca spp. all to 40ft. Acacia spp. sp. 3, DCantherhania concolor, Hypochoeris sp. 1, to 20ft + regrowth, Pandanus sp. to 15ft. Spenbeckantsp. Vitripennis, S. Rieu, Caledia Grasses mixed 1-4ft, occ. forbs. Also Captiva, Aciolopus thalassinus, Adperrisia acola, a former gravel dump with short grass Abroctonospha similis, Bundinga sp. P70, Spectriforme + bare gravelly ground. Gastr., Austr. basal, transversa Caledia, Acida, Austr. gutt.

11110.6 Back on bitumen. Tunes R.

11111.1. Tunes R.

11112.2 End of road at Golf Club. Beige sand hills, very jumbled. Sparse grasses

Sorghum stipoideum (Ewart & White) Gardn. & Hubb. 66
(spec. ?: moorabinensis on this)
to 2ft ½. pink forb to 1ft. bare sand.
Genus nov. 70 sp. 1, Lyperobia sp. 1, Xanthorrhiza
Patches of mixed broad-leafed shrubs &
concolor, Pycnostictus seriatus, Acacia Conica,
small trees, occ. Acacia & oblong-leafed
Adelappa erythroptera, Hastella sp. P208, Spectriforma transversa
occ. at landward margins. Return.

11116.3. Hotel.

11118.2 Airport intersection. Turned R.

11119.7. Road off to L.

11124.6. Water on L.

11126.7. Cross conveyor belt.

11128.9. End of road at treatment plant.
Return.

11131.1. Turned R.

11135.9. Drimmie Hill & adjacent sandflat.
Goniacordea sp. 3, Tolgadea, inferma, Genus
The hill has massive granite outcropping
nov. 70 sp. 1, Genus nov. 98 sp. 1, Abaitoniopsha
& bears monsoon forest to 30ft & open
similis, Daperria actola, Spectriforma transversa,
patches of grass to 6ft. Lower area
Hastella sp. P208
with grass & forbs to 2½ft under occ.
Casuarina & occ. Pandanus. Cockroaches
& earwigs in Pandanus. Moorabinensis mainly
on a dense patch of leguminous forbs to
1ft on sand flat. Return.

19/5.

111.51.2 Hotel.

179.9. Dhupuma. Tarned R.

187.1. Flat with beige sand & sparse.
Genus nov. 98 sp. 1, Genus nov. 105 sp.,
laterite nodules on underlying massive
Restibropis sp.; Canterraria concolor, Genus nov.
laterite. E. tetrodonta, fairly dense, to
70 sp-1, Dappera arcola, Gonioalcoca sp. 3,
60ft, with Livistona dense to 8ft, occ.
Genus nov. 61 sp. 2, Macrotona sp. 12, Gonioalca
Banksia sp. & Grevillea sp. to 8ft, a
vocans, Acacia gonocarpa F. Muell. Dipodium
very fine-phyllode Acacia to 1ft abundant,
Hibbertia dealbata (DC.) Benth., Hibbertia lepidota DC.
Hibbertia spp. (2) to 1ft, other forbs occ.,
sp-1, Bundinja sp. P76, Spectrifomia transversa
grasses 1–6ft frequent, v. occ. Calytrix
to 4ft.

190.5. Slopes with a complete cover of
Gonioalca vocans, Macrocarpa conglobata
coarse laterite gravel & occ. rocks.
Gonioalca sp. 3, Canterraria concolor,
E. tetrodonta to 60ft fairly dense, with
Restibropis sp., Bundinja sp. P76
Livistona to 5ft, Acacia sp. to 10ft,
Acacia sp. with small phyllodes (1cm) to
4ft, Hibbertia dealbata (DC.), H. lepidota DC.
Hibbertia spp., taller small shrubs
& forbs, scattered grasses 1–4ft, some
bare ground.

194.7 First crossing, Giddy R.

194.9 Road off to L.

195.1. Main Eddy crossing.

198.8 Track off to L.

210.6. Slight slope with brown soil densely covered with laterite pebbles. E. tetrodonta fairly dense to 50ft, with occ. Erythrophleum to 25ft, occ. sessile-leafed Eucalyptus setosa Schau. angustifrons, Acacia woodwood (occ.) to 15ft, Livistona to 5ft abundant, occ. Pandanus regrowth, broad-phyllode Acacia to 6ft (marabinna on this), Hibbertia spp. to 1ft, other forbs occ, Maclyptria on short grass. mixed grasses 9in to 6ft. Return.

211.2. Edge of scarp. Undulating stony laterite. E. tetrodonta to 50ft fairly dense, with Livistona abundant to 5ft, Acacia spp. occ., grasses sparse to 1–5ft, forbs occ., Erythrophleum regrowth occ.

241.0 Obuseima.

26/5.

11299.5 Hotel.

301.4 Tunes L.

302.5. Gentle depression with pale sand. Melaleuca spp. to 20ft, occ. E. tetrodonta

Genus nov. 70 sp. 1, Xylopia sp. 1, Macrotona sp. 12(?) (j), Xanthorrhiza concolor, Heteropteris Obscura, Gonioalvocans, + Acacia spp. Dapsilia accola, to 40ft, Grevillia sp. to 12ft, Pandanus Aciotopis thalassinus, Tolgadia bivittata, at lowest point to 12ft patches round- Genus nov. 98 sp. 1, Calycia capitata, leafed Kurrajong, patches Calytrix, grassy Stenocarpus sinuatus, Heteropteris, Rechtorpis sp. 1—4ft, patches open sedge to 10in, occ. Hibbertia decalvata (DC.) Benth. Spectriforma transversa, large-leafed Hibbertia. Return.

Hastella sp. P208
Bundinja sp. P70

305.7. Hotel.

307.5. Turned R.

309.2. Gentle undulations on pale sand. Erythropoumala amaena, Goniocoidea sp. 3, occ. with a little lateritic gravel. E. Macrotona sp. 12(?)(j.), Xanthorrhiza concolor, tetrodonta to 60ft open, with Grevillea + in places Melaleuca spp. Rechtorpis sp. sp. to 15ft, occ. Banksia spp. to 8ft. Acacia Genus nov. 98 sp. 1, Genus nov. 105 sp. sp. a other shrubs to 5ft, Hibbertia spp. Spectriforma transversa, Bundinja sp. P70 (3) to 2ft, scattered grasses + sedges to 18in, occ. Livistona to 3ft + occ. Pandanus regrowth, much litter + nymphs some have grazed. Genus nov. 106 for Hibbertia lepidota DC. finer leafed Hibbertia from 11024.8.

Gonioalvocans

21/5.

11311.8 Hotel.

3440.7. Dhupuma.

3441.9. Turned L.

Site (6)

11353.4 Near end of road opposite ... all inland. Sandhills + granite + ... Aristida, Aciotis thalassoides, Kanferia, Heterosperma obscurella, Calellia ... Low spreading ... (2) to 1ft, occ. Pandanus to 6ft, occ. oblong-leaf ... Acacia sp. 12 (3), Grevillea ... to 6ft, patches of coastal scrub, ... Spectriforma transversa grasses to 4ft + low annual grasses to 6in; some forbs + ... gadea sp. 1, Stylobasium ... Site 7. Estuarine type of mulch from Dalyellup, Adelppa erythrobotrya, Heterosperma obscurella, ... swamp with dark mud + water, Lobelia concolor, ... Pachistopetalum sp. 1 ... grasses to 12ft + ... coarse grasses to 5ft scattered

Site 8. Low hills + sandhills with ... Acacia sp. 12 (3), Grevillea ... patches of scrub + scrub ... to 2ft + some other grasses, Heterosperma ... with pink flower. Some sand.

Returns.

11354.6 Swampy flat with grey mud. ... Acacia sp. 1, Eucalyptus sp. 1, Xanthorrhoea ... Melaleuca sp., all to 15ft, Spectriforma transversa ... grasses to 1ft, occ. ...

3400.1 Stony ... slope ... to

[illegible] E. diffusa to to
60ft., Cunninghamia to 15ft. Acacia sp. rare
to 20ft., Eucalyptus sp. (2) to 40ft. Hibbertia
sp. to 2ft + other forbs + shrubs incl.
Epacrids, Grevillea. Acacia prasinus.

366.6. Dracophyllum.

32/5.

11395.3. Hotel. Track + port roads.

405.1. Tussock L.

414.2 = 405.1. (more silky wet.)

442.1 2mile NW of Gem Creek. Apex of
NW-directed promontory. Slightly swampy
flat with Melaleuca spp. (2). oblong-elliptical
trees (2). occ. E. papuana, occ. E. ptychocarpa
Gaultheria, Tolyadium, Epacris, Cyathodes, Dracophyllum
+ E. minuta on fringes. Grasses dense
to 4ft + patches to 1ft. occ. Pandanus
to 7ft. occ. shrubs. incl. Rumohra.

446.3 Swamps with dark mud, lined
with Melaleuca spp + bloodwoods to
30ft, heath to 12ft. mainly Epacrids
Gaultheria, Cyathodes, Dracophyllum, Leptospermum
Thelionema, Astelia, Gleichenia, Lycopodium
Restio to 12ft + thick grasses + forbs

[illegible] with [illegible] spectiformis [illegible]
[illegible] with low dense bearing Lagunata at low density with
[illegible] rotundata [illegible]

4.56.0 (Index exceeded) Main intersection

Track road to treatment plant

4.62.5 Turned round.

4.63.5 Margin of inlet with mangroves Hastella sp. P208
Tolypella, Nitella, Macrocystis, Cauloglossum, Enteromorpha, Grasses & sedges to 6 in. to 1 ft.
E. hispanica, Halodule, Thalassia, Syringodium to 4 ft. wide, big with ball 7 ft. Thalassia
dominant over-grazing under E. hispanica & Halodule. Thalassia
& Halodule. Thalassia

4.69.0 Main intersection.

23/5

11.34.7 Hotel

4.16.6 Turned to [illegible]

4.16.0 Quarter

4.14.6 Flat with [illegible] E. hispanica to
Thalassia, Halodule, Cymodocea, Nitella, Chara sp. &
E. hispanica to 6 ft. with [illegible]
Acetabularia sp. & other small [illegible]

Mangrove grasses 1 – 5 ft.

5.05.2 Back on main road.

Vanuatu. The whole area is tectonically active. The uplift has been considerable to explain the underlying granitic limestone on the east coast, on Brumiel Peninsula, and occasionally elsewhere), in the swampy swampy plains (the largest behind Vanuatu), and in the deformation controlled complexes of the north and south-east coastlines. The only area of sandstone was a small area, associated with the granites, constituting the hills and headlands that alternate with mangroves on the east coast south of Lautoka Bay. The area was mostly granitic and hilly, impenetrable on the way of flora and swamp on the way of grassy hillsides. It is the absence of sandstone (or granite, or conglomerate), and with it of Triodia (except for some

possible occurrence in open
sea plankton in Antarctic waters).
Next must be considered the
responsible for the large difference in
the grasshopper faunas of western
Argentina and the Gaspé Peninsula,
although the more marked climatic
of the latter may also play a part.

The grassy plain near Hinton
was unfortunately not visited, but it
is unlikely to be radically different
from similar localities near the
[illegible] Hypochaetis floridana
continued beneath the edge of the
coastal sand covering its host
immediately behind the narrow
lagoon bordering the sandy ridge
which [illegible] the Gulf. The
grass in this region is [illegible]
in tropical Southern Florida, but the
plants are smaller. No Eritettix obtusa
were seen, but an extensive

search was not made.

Valanga nubleger & the genus Valanga sp. were found only in the coastal scrub behind the hotel. At first it was thought this might be due to our visiting this site early in the morning, when the insects were basking on the outer foliage. However, investigation showed that the number (without collecting) seen on a selected stretch of the scrub increased from 19 at ca. 8 am to 25 at ca. 11 am., & subjective observation suggested that even more could have been counted a couple of hours later. It was possible to collect some 30–50% of those seen, & to repeat this harvest with the same success three or four times at about hourly intervals. Apparently many individuals are deeper in the scrub at any one time. This seemed to be a preference for the lower

5ft or so, + indeed for (of 1-2 m height throughout) a couple of metres in front of the main face. Some were on the sand among grass in front of the face — apparently more females than males. V. irregularis was also common than the other two, but was seen occasionally at other sites.

Both smaller species were infested with green tracheal mites, which sometimes virtually filled the air sacs + major tracheae, along with numerous translucent mite eggs. Infestation was much heavier in males. Two or three recently dead individuals were found. It seems possible that the Valanga populations are subject to drastic crashes due to this parasite, + can build up only in widely separated patches. No particularly sluggish individuals were found,

however. The possibility that the population had been started from eggs introduced with garden plants for the hotel has to be considered. Although none were found in the garden (one *irregularis* was), the latter is sprayed every few days with some insecticide, presumably for mosquito control. The only unquestionably natural occurrence of *meleagris* ♀ of a green Valanga (same species? — taxonomy unsatisfactory) known to me from personal observation is in [deleted] the fringes of] a disturbed monsoon forest on Mt. White, at Coen. The Kimberley records were presumably associated with cultivation.

The three species of *Talgadia* showed increasing wet-environment preference in the order *infirma*, *hewitsata*, sp. 1. *Oxya japonica* seemed to have a similar preference to *T.* sp. 1, but usually on shorter grass. *Daprova*

Hosts found on Lipari Island. Locality of Brandaris, from which to collect by dredging & hauling.

Two species represented by many single specimens, more upon [illegible] specimens & more still upon single localities. All these mollusks were taken at many localities, two of them in considerable numbers. It is unlikely that any species escaped detection.

The life-cycle of Teredo navalis L. (W. Rathbun)
is curious & worth examining. Larvae in November & May (W. Rathbun & Stone), young adults in March (W. A. Herdman). Specific differences may possibly lie in [illegible]